



CONSENT FOR KYBELLA INJECTION

Kybella is an FDA approved cosmetic injection indicated for improvement in the appearance of sub mental fullness associated with submental fat in adults.

The results of Kybella are not immediate. At your first treatment visit you will receive a series of injections in the submental area. At the time of injection and for several days following the injections you will have swelling in the submental area. Kybella will cause the fat cells to diminish gradually over the course of the next month following your injection in the treated area. A series of treatments may be necessary to achieve optimal results and these will occur at no sooner than the one month interval. Your doctor will decide the appropriate number of treatment sessions and the amount of Kybella you will need at each session.

Side Effects:

Side effects of Kybella may include: bruising, swelling, numbness, induration, marginal mandibular nerve injury, dysphagia (difficulty swallowing), bleeding, tenderness or discomfort, tissue necrosis, hyperpigmentation, redness, or alopecia at the site of injection.

Alopecia: Kybella injections could cause small patches of alopecia in the treatment area.

Pre-Instructions: You should not be pregnant, nursing an infant, have a history of a bleeding disorder or infection of the area to be treated. The morning of your treatment you should take ibuprofen if you are able to tolerate this medication. Also, you should tell you physician if you take any blood thinners or aspirin or have a history of any issue with lidocaine or epinephrine.

Post Care Instructions:

- Ice packs may be used to the treated area during the first 12 hours – ice for 15 minutes on every hour • Do not massage the injection site
- If you are able to tolerate ibuprofen take 200mg three times daily beginning the day of your treatment and for the following four days
- Avoid heavy exercise the day of your treatment
- Notify Alladerm if any significant, swelling, bleeding, pain, dusky discoloration, difficulty swallowing or smiling, or fever occurs.

Alternatives: As explained not all submental fullness will respond to Kybella. Other alternative treatments are liposuction to the area, a neck lift and/or platysmoplasty procedures.

I certify that I have read the above consent and fully understand it. I have been given ample opportunity for discussion and all my questions have been answered to my satisfaction. I hereby consent to the Kybella injection procedure. This constitutes the full disclosure and supersedes any previous verbal or written disclosures.

Signature-Patient or Guardian

Print Name/Relationship

Date

Signature-Witness

Print Name

Date